

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07H 21/02, 21/04, C07K 9/00 A61K 37/00, C12N 1/21, 5/10 C12N 15/70, C12P 21/00		A1	(11) International Publication Number: WO 94/10187 (43) International Publication Date: 11 May 1994 (11.05.94)
(21) International Application Number: PCT/US93/10307 (22) International Filing Date: 26 October 1993 (26.10.93) (30) Priority data: 07/968,953 30 October 1992 (30.10.92) US 07/968,954 30 October 1992 (30.10.92) US (60) Parent Applications or Grants (63) Related by Continuation US 07/968,953 (CIP) Filed on 30 October 1992 (30.10.92) US 07/968,954 (CIP) Filed on 30 October 1992 (30.10.92) (71) Applicants (for all designated States except US): HSC RESEARCH AND DEVELOPMENT LIMITED PARTNERSHIP [CA/CA]; 88 Elm Street, Toronto, Ontario M5G 1X8 (CA). MEMORIAL SLOAN-KETTERING CANCER CENTER [US/US]; 1275 York Avenue, New York, NY 10021 (US). CONSEJO SUPERIOR DE INVESTIGACIONES CIENTIFICAS [ES/ES]; Velazquez, 144, E-28006 Madrid (ES).		(72) Inventors; and (75) Inventors/Applicants (for US only): LETARTE, Michelle [CA/CA]; 35 Howland Road, Toronto, Ontario M4K 2Z7 (CA). MASSAGUE, Joan [US/US]; 1233 York Avenue, New York, NY 10021 (US). BERNABEU, Carmelo [ES/ES]; Centro Investigaciones Biologicas, Velazquez, 144, E-28006 Madrid (ES). CHEIFETZ, Sela [CA/CA]; 88 New Seabury Drive, Concord, Ontario L4K 2R8 (CA). (74) Agents: KONSKI, Antoinette, F. et al.; Campbell & Flores, 4370 La Jolla Village Drive, Suite 700, San Diego, CA 92122 (US). (81) Designated States: AU, BB, BG, BR, CA, CZ, FI, HU, JP, KP, KR, LK, LV, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, US, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
Published with international search report AMIL INFORMATION SERVICES PO BOX 405, CORTE MADERA, CA 94976-0405 (415) 927-0340 • FAX (415) 927-7250			
(54) Title: COMPOSITIONS AND METHODS FOR MODIFYING THE REGULATORY ACTIVITY OF TGF- β			
(57) Abstract <p>This invention provides a novel purified TGF-β binding glycoprotein, endoglin, and isolated nucleic acid molecules that encode amino acid sequences corresponding to the TGF-β-binding glycoprotein. Also provided is soluble endoglin-derived polypeptide, and fragments thereof. A pharmaceutical composition which comprises the purified endoglin-derived polypeptide or produced recombinantly methods and a pharmaceutically acceptable carrier is further provided as well as methods of treating patients by administering to the patient the pharmaceutical compositions of this invention.</p>			

BEST AVAILABLE COPY

62A1
ATTORNEY DOCKET NUMBER: 10177-216-999
SERIAL NUMBER: 09/590,002
REFERENCE: B107

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07H 21/02, 21/04, C07K 9/00 A61K 37/00, C12N 1/21, 5/10 C12N 15/70, C12P 21/00		A1	(11) International Publication Number: WO 94/10187 (43) International Publication Date: 11 May 1994 (11.05.94)
(21) International Application Number: PCT/US93/10307 (22) International Filing Date: 26 October 1993 (26.10.93) (30) Priority data: 07/968,953 30 October 1992 (30.10.92) US 07/968,954 30 October 1992 (30.10.92) US (60) Parent Applications or Grants (63) Related by Continuation US 07/968,953 (CIP) Filed on 30 October 1992 (30.10.92) US 07/968,954 (CIP) Filed on 30 October 1992 (30.10.92) (71) Applicants (for all designated States except US): HSC RESEARCH AND DEVELOPMENT LIMITED PARTNERSHIP [CA/CA]; 88 Elm Street, Toronto, Ontario M5G 1X8 (CA). MEMORIAL SLOAN-KETTERING CANCER CENTER [US/US]; 1275 York Avenue, New York, NY 10021 (US). CONSEJO SUPERIOR DE INVESTIGACIONES CIENTIFICAS [ES/ES]; Velazquez, 144, E-28006 Madrid (ES).		(72) Inventors; and (75) Inventors/Applicants (for US only) : LETARTE, Michelle [CA/CA]; 35 Howland Road, Toronto, Ontario M4K 2Z7 (CA). MASSAGUE, Joan [US/US]; 1233 York Avenue, New York, NY 10021 (US). BERNABEU, Carmelo [ES/ES]; Centro Investigaciones Biologicas, Velazquez, 144, E-28006 Madrid (ES). CHEIFETZ, Sela [CA/CA]; 88 New Seabury Drive, Concord, Ontario L4K 2R8 (CA). (74) Agents: KONSKI, Antoinette, F. et al.; Campbell & Flores, 4370 La Jolla Village Drive, Suite 700, San Diego, CA 92122 (US). (81) Designated States: AU, BB, BG, BR, CA, CZ, FI, HU, JP, KP, KR, LK, LV, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, US, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
		Publ. Int. Information search report AMI INFORMATION SERVICES P O BOX 405, CORTE MADERA, CA 94976-0405 (415) 927-0340 • FAX (415) 927-7250	
(54) Title: COMPOSITIONS AND METHODS FOR MODIFYING THE REGULATORY ACTIVITY OF TGF- β			
(57) Abstract <p>This invention provides a novel purified TGF-β binding glycoprotein, endoglin, and isolated nucleic acid molecules that encode amino acid sequences corresponding to the TGF-β-binding glycoprotein. Also provided is soluble endoglin-derived polypeptide, and fragments thereof. A pharmaceutical composition which comprises the purified endoglin-derived polypeptide or produced recombinantly methods and a pharmaceutically acceptable carrier is further provided as well as methods of treating patients by administering to the patient the pharmaceutical compositions of this invention.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

COMPOSITIONS AND METHODS FOR MODIFYING THE REGULATORY
ACTIVITY OF TGF- β

FIELD OF THE INVENTION

The present invention relates to cell biology and
5 to methods of modifying the biological activity of cell
regulatory factors. More specifically, the present
invention relates to a novel TGF- β -binding glycoprotein.

Throughout this application various publications
are referenced within parentheses. The disclosures of
10 these publications in their entireties are hereby
incorporated by reference to more fully describe the state
of the art to which this invention pertains.

BACKGROUND OF THE INVENTION

Glycoproteins, in which one or more carbohydrate
15 units have been attached covalently to the protein by
posttranslational processing are widely distributed.
Several secretory proteins, including the immunoglobulins,
are glycoproteins, as are most components of plasma
membranes such as cell membrane receptors, where the
20 carbohydrates can be involved in cell-to-cell adhesion.

Transforming growth factor β (TGF- β) refers to a
family of multi-functional cell regulatory factors produced
in various forms by many cell types (for review see Sporn
et al., J. Cell Biol. 105:1039 (1987)). Five distinct
25 isoforms of TGF- β have been identified. TGF- β 1 and TGF- β 2
have been characterized in detail. TGF- β is the subject of
U.S. Patent Nos. 4,863,899; 4,816,561 and 4,742,003 which
are incorporated herein by reference. TGF- β binds to cell
surface receptors present on various types of cells and is
30 known to potentiate or inhibit the response of most cells
to other growth factors, depending on the cell type. TGF- β
also regulates differentiation of some cell types, either
promoting or inhibiting proliferation of the cell. Another

marked effect of TGF- β is the promotion of cellular production of extracellular matrix proteins and their receptors (for a review see Keski-Oja et al., J. Cell Biochem. 33:95 (1987); Massague, Cell 49:437 (1987);
5 Roberts and Sporn, "Peptides Growth Factors and Their Receptors", Springer-Verlag (1989)).

Notwithstanding the beneficial and essential cell regulatory functions served, TGF- β regulatory activity can prove detrimental to its host organism. For example,
10 whereas growth and proliferation of mesenchymal cells is stimulated by TGF- β , some tumor cells may also be stimulated, using TGF- β as an autocrine growth factor. In other cases the inhibition of cell proliferation by TGF- β similarly proves detrimental to its host organism. An
15 example would be the prevention of new cell growth to assist in repair of tissue damage. The stimulation of extracellular matrix production by TGF- β is essential for wound healing. However, in some cases, the TGF- β response is uncontrolled and an excessive accumulation of
20 extracellular matrix results. An example of excessive accumulation of extracellular matrix is the "internal" scarring that occurs in the pathology glomerulonephritis and dermal scar tissue formation.

25 The transforming growth factor- β receptor system in most mesenchymal and epithelial cells consists of several components (Massague, J. Ann. Rev. Cell Biol. 6:597 (1990); Lin, H.Y. et al., Cell 68:775 (1992); Georgi, L.L. et al., Cell 61:635 (1990); Mathews, L.S. et al., Cell
30 65:973 (1991); Attisano, L. et al., Cell 68:97 (1992); Lopez-Casillas et al., Cell 67:785 (1991) and Wang et al., Cell 67:796 (1991)), one of which is betaglycan, a membrane-anchored proteoglycan. In addition to betaglycan, the TGF- β receptor system in most mesenchymal and
35 epithelial cells consists of the type I receptor, a 53-kDa glycoprotein whose structure has not been determined yet,

and the type II receptor, which belongs to the protein serine/threonine kinase receptor family. Additional cell surface TGF- β -binding proteins, some of which have a more restricted distribution, have also been described.

- 5 Thus, a need exists to develop compounds that can modify the effects of cell regulatory factors such as TGF- β . The present invention satisfies this need and provides related advantages as well.

SUMMARY OF THE INVENTION

- 10 The present invention provides a novel purified TGF- β -binding glycoprotein. The protein, endoglin, is expressed at high levels on human vascular endothelial cells.

- Further provided by the present invention are
15 methods of treating pathologic conditions mediated by TGF- β regulatory activity by contacting the TGF- β with an effective amount of purified endoglin-derived polypeptide or any fragment thereof having the ability to bind TGF- β . Thus, intact, native endoglin and soluble fragments thereof
20 are useful in these methods. This invention provides a method of preparing and purifying full length and soluble endoglin-derived polypeptide. Isolated nucleic acids encoding the novel TGF- β -binding glycoprotein and soluble endoglin-derived polypeptides are also provided, as well as
25 vectors containing the nucleic acids and recombinant host cells transformed with such vectors.

BRIEF DESCRIPTION OF THE FIGURES

- Figure 1 shows the domain structures of betaglycan and endoglin. Shown is a schematic
30 representation highlighting regions of similarity between the linear sequences of betaglycan, an 853-amino acid

transmembrane proteoglycan, and endoglin, a disulfide-linked transmembrane protein composed of two identical subunits of 633 amino acids each. The transmembrane and short cytoplasmic regions (dark shaded box) of endoglin have a high level of sequence similarity to the corresponding regions of betaglycan. Two regions of weaker similarity are detected in the ectodomains of these proteins (light shaded boxes). Numbers represent the percent amino acid sequence similarity between the indicated domains of betaglycan and endoglin. Closed ovals represent positions of cysteine residues. Two putative sites for glycosaminoglycan chain attachment in betaglycan are indicated.

Figure 2 shows cell surface TGF- β 1-binding proteins expressed by HUVEC. Near confluent cultures of HUVECs were affinity-labeled by incubation with 100 pM ^{125}I -TGF- β 1 followed by chemical cross-linking with 0.16 mM disuccinimidyl suberate. A) Triton X-100 extracts of affinity-labeled HUVEC were resolved on SDS-PAGE gels under reducing (R) or nonreducing (NR) conditions. Lane C contains extract from cells affinity-labeled in the presence of excess unlabeled TGF- β 1. The migration position of TGF- β receptors I (RI) and II (RII) are indicated. Arrow, the major affinity-labeled proteins of 180 kDa and higher molecular mass apparent on nonreducing gels. Arrowhead, the affinity-labeled proteins of 110-120 kDa seen on reducing gels. B) Detergent extracts of affinity-labeled HUVEC were resolved under nonreducing conditions on a first gel that was then resolved under reducing conditions in the second dimension as described in Cheifetz and Massague, *J. Biol. Chem.* 266:20767-20772 (1991), incorporated herein by reference. The 110-120-kDa labeled species migrating off-the-diagonal are indicated (arrowheads).

Figure 3 shows specific immunoprecipitation of TGF- β 1-endoglin complexes. HUVECs were affinity-labeled with 100 pM 125 I-TGF- β 1 as described in Figure 2. A) Detergent extracts of affinity-labeled cells were incubated with mAb 44G4 and immune complexes were collected on protein G-Sepharose. After washes, equal aliquots of the samples were analyzed under reducing (R) or non-reducing (NR) conditions by SDS-PAGE (5-8% polyacrylamide gradient gels). B) Affinity-labeled HUVEC lysates were maximally depleted of endoglin by two successive 45 min incubations at 4°C with 100 μ l of 44G4-IgG-Sepharose. S) supernatant after second immunoprecipitation. I) the first 44G4 immunoprecipitation which contained 83% of the endoglin. T) corresponding amount of total extract used for the depletion experiment. All samples were analyzed under nonreducing conditions on SDS-PAGE with the exception of I_R, which was run under reducing conditions. The migration positions of TGF- β receptor II (RII), and endoglin monomer, dimer, and oligomer are indicated.

Figure 4 shows that endoglin transiently expressed in COS-M6 cells binds TGF- β 1. COS-M6 cells were transfected with a cDNA encoding full-length L-endoglin (Endoglin) or control vector (C). Cells were affinity-labeled with 150 pM 125 I-TGF- β 1 and the detergent extracts incubated with mAb 44G4 followed by protein G-Sepharose. Immunoprecipitated proteins were analyzed by SDS-PAGE under reducing (R) and non-reducing (NR) conditions and visualized by autoradiography.

Figure 5 shows the specificity of endoglin for TGF- β isoforms assessed in COS cell transfectants and in HUVEC. A) COS-M6 cells transfected with endoglin vector were affinity-labeled with 150 pM 125 I-TGF- β 1 alone or in the presence of 1 or 10 nM unlabeled TGF- β 1, - β 2 or - β 3. B) HUVEC were affinity-labeled with 100 pM 125 I-TGF- β 1 alone or in the presence of 5 nM unlabeled TGF- β 1 or TGF- β 2.

Lysates from these cells were immunoprecipitated with MAb 44G4. Immunoprecipitates were fractionated under reducing conditions on SDS-PAGE gels. The region of the gels containing monomeric endoglin is shown along with the migration position of 100-kDa marker.

Figure 6 shows the restriction enzyme map of vector pcNeoSolEND with the 1.7 kb endoglin cDNA insert.

Figure 7 shows the partial nucleotide and predicted amino acid sequences of isolated S-endoglin cDNA. Nucleotides are numbered on the right. Amino acids are numbered on the left. The predicted signal sequence (nucleotides 283-357) and transmembrane region (nucleotides 2042-2116) are boldfaced. The 135 bp insert (underlined) contains splicing consensus sequences of donor/acceptor sites (GT, AG at positions 2134 and 2267) and branch point of lariat (CTGAC at position 2234). Nucleotides 372-2021 and 2322-3073 were found to be identical to the corresponding cDNA sequence of endothelial endoglin (Gougos, A. and Letarte, M.J., J. Biol. Chem. 265:8361 (1990) and Table 1).

Figure 8 shows the analysis of the cytoplasmic region which reveals the existence of two different forms of endoglin. A. Diagram of the cDNA coding for the cytoplasmic regions of the two alternative forms of endoglin. Only the region corresponding to the 3' end containing the cytoplasmic domain is depicted. The isolated endoglin cDNA (S-endoglin) contains a 135 bp insert not present in the previously described sequence (L-endoglin) (Gougos, A. and Letarte, M.J., J. Biol. Chem. 265:8361 (1990)). The sequence of the additional 135 bp insert is shown in Figure 7. The position of the stop codons and the corresponding translated protein sequences (thick bar), are indicated. B. Alignment of the cytoplasmic and transmembrane domains of S-endoglin with

the corresponding sequences of L-endoglin and human betaglycan, Morén, A. et al., Biochem. Biophys. Res. Commun. 189:356 (1992). Boxes contain identical sequences. Numbers indicate the position of the first amino acid in the whole sequence. Two major regions of identity were found. The first region (73% identity) involves residues 587-617 of S- and L-endoglin and residues 780-810 of betaglycan. The second region (74% identity), involves residues 634-660 of L-endoglin and residues 823-849 of betaglycan. The transmembrane region of S-endoglin is boldfaced. Dashes have been inserted for purposes of alignment. Asterisks indicate the last residue of the protein.

Figure 9 shows the expression of L-endoglin and S-endoglin in transfectant cells. Mouse fibroblasts were transfected with either L-endoglin or S-endoglin cDNA and the expression of the endoglin molecule analyzed. A. Analysis by cytofluorometry of the endoglin present at the cell surface. After trypsinization, cells were stained for indirect immunofluorescence with the monoclonal antibody 8E11 (anti-endoglin). A control staining of endoglin mock transfectants is also shown. B. Immunoprecipitation analysis. Cells were metabolically labeled with [³⁵S]methionine, lysed and immunoprecipitated with 44G4 (anti-endoglin) or HC1/1 (anti CD11c) monoclonal antibody. Monoclonal antibody HC1/1 was included as a negative control. Samples were electrophoresed on a 6-12% acrylamide gradient gel under nonreducing conditions. C. Immunoblotting analysis. Mock (L cells), L-endoglin (L-Endo) and S-endoglin (S-Endo) cDNA transfected mouse fibroblasts and PMA-treated U937 cells were lysed in Triton X-100, and the insoluble material removed by centrifugation. Proteins contained in the supernatant were electrophoresed on a 6% acrylamide gel under non-reducing conditions and transferred to nitrocellulose membranes. Immunodetection of endoglin was carried out with 44G4

(anti-endoglin) monoclonal antibody using a chemiluminiscent assay. HCl/1 (anti-CD11c) and X63 monoclonal antibody were used as negative controls. D. Immunoprecipitation analysis of cell surface labeled endoglin. Mock, L-endoglin (L-Endo) and S-endoglin (S-Endo) transfected mouse fibroblasts were ¹²⁵I-labeled, lysed and immunoprecipitated with 44G4 monoclonal antibody (anti-endoglin). Samples were electrophoresed on a 10% acrylamide gel under either reducing (R) or nonreducing conditions (NR).

Figure 10 shows the detection of L-endoglin and S-endoglin transcripts by PCR amplification. Samples of total RNA from placenta, PMA treated HL-60 cells or PMA-treated U937 cells were incubated either in the presence (+RT) or in the absence (-RT) of reverse transcriptase. The generated cDNA samples, together with cDNA from S-endoglin (S-Endo) or L-endoglin (L-Endo) clones in pUC13 and cDNA from an endothelial cell library, were used for PCR amplification in the presence of oligonucleotides #14 and #15 specific for S-endoglin (panel A), or oligonucleotides #12 and #11 common to both L-endoglin and S-endoglin (panel B). No amplification was observed when the RT reaction was omitted, excluding the possibility of DNA contaminating the RNA samples. Additional bands below the specific amplified fragments of L and S endoglin probably represent primer-dimer artifacts.

Figure 11 shows that both S-endoglin and L-endoglin bind TGF-β1. Confluent cultures of S-endoglin (L+S) and L-endoglin (L+L) transfectants were affinity labeled by incubation with 100 pM ¹²⁵I-TGF-β1 alone or in the presence of 4 nM unlabeled TGF-β1, followed by chemical crosslinking with disuccinimidyl suberate. All samples bound specifically to TGF-β1, as revealed by a fourfold ratio between the cpm bound in the absence versus presence of cold competing ligand; the parental L cells and mock

transfectants bound on average 60,000 specific cpm, while the endoglin transfectants bound on average 110,000 specific cpm. Immunoprecipitates of the L+-S transfectant with 44G4-IgG Sepharose contained 7700 cpm on average
5 versus 700 cpm with control IgG-Sepharose; immunoprecipitates of the L+-L transfectant contained 4300 cpm versus 630 cpm for control. These immunoprecipitates were run on a 6-9% acrylamide gradient gel in reducing (R) or nonreducing (NR) conditions. The positions of the endoglin
10 monomer, dimer and oligomer and of molecular weight markers are indicated.

Figure 12 shows the multiple cloning sites of pcDNAI/Neo.

Figure 13 shows the amino acid and nucleotide
15 sequence of "L-Endoglin."

Table 1

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
84

DETAILED DESCRIPTION OF THE INVENTION

Endoglin is a homodimeric membrane glycoprotein composed of disulfide-linked subunits. Human-derived endoglin has been shown to exist in at least two isoforms expressed from a human cDNA library, an "L-isoform," of about 90 kDa or an "S-isoform" of about 85 kDa, reduced. Human endoglin, purified from tissue is shown to be composed of two disulfide-linked subunits each of about 95 kDa. It is expressed in human pre-erythroblasts, macrophages, leukemic cells of the lymphoid and myeloid lineages and at higher levels in vascular endothelial cells. It is also abundant on the syncytiotrophoblast, the multinucleated placental layer which constitutes the interface with maternal blood and plays an important role in providing nutrient exchange and immunological protection of the fetus (Gougos et al., Inter. Immunol. 4:83-92 (1992)).

Endoglin was first identified on a pre-B leukemic cell line, by its reactivity with mAb 44G4 (Quackenbush and Letarte, J. Immunol. 134:1276-1285 (1985)). It is present at low levels on cells derived from childhood acute leukemia cases (ALL) with pre-B lymphoid and myeloid phenotype; it is absent from T-ALL (Gougos and Letarte, J. Immunol. 141:1925-1933 (1988b); Kreindler et al., Leukemia and Lymphoma 3:7-18 (1990)). Interestingly, the human endoglin gene is localized to chromosome 9q34-qter, likely within the region translocated to chromosome 22 in Philadelphia chromosome-positive leukemia (Fernandez-Ruiz et al., Cytogen. Cell Gen. 64:204-207 (1993)). In normal adult bone marrow, only 3-5% of mononuclear cells express endoglin and they bear a pro-erythroblast phenotype; pre-B and myeloid precursors do not show detectable levels of endoglin (Buhring et al., Leukemia 5:841-847 (1991)). Normal B and T lymphocytes and unstimulated monocytes do not express endoglin but an up-regulation is observed on

activated macrophages (Lastres et al., Eur. J. Immunol. 22:393-397 (1992)).

Endoglin expression is considerably increased in the endothelium of various pathological skin lesions where endothelial cell proliferation is known to occur (Westphal et al., J. Invest. Dermatol. 100:27-34 (1993)). In tumors, capillary endothelial cells undergoing active angiogenesis also show higher levels of endoglin than resting endothelium in adjacent tissue.

Purified human endoglin exists in various isoforms and is composed of two disulfide linked subunits of Mr=95,000; 90,000; or 85,000, and bears N- and O-linked oligosaccharides. The primary sequence of human endoglin is composed of a 25 amino acid signal sequence, an extracellular domain of about 561 amino acids, a single transmembrane region of 25 amino acids and a cytoplasmic tail of 14 to 47 amino acid residues (Gougos and Letarte, J. Biol. Chem. 265:8361-8364 (1990)).

A relationship between human endoglin and the TGF- β receptor system was discovered with the molecular cloning of the rat TGF- β -binding proteoglycan, betaglycan (also known as the type III TGF- β receptor), which revealed that the transmembrane domain and the relatively short (43 amino acid) cytoplasmic tail of this protein were remarkably similar (71% amino acid sequence similarity and 63% amino acid identity) to the corresponding regions in endoglin (see Figure 1). The extracellular domains of these two proteins show limited homology in primary structure, and while endoglin is not a proteoglycan, it does contain N- and O-linked oligosaccharides.

Cloning of the TGF- β receptor II (Lin et al., Cell 68:775-785 (1992)) revealed a functional transmembrane serine/threonine kinase which had previously been

identified as a polypeptide of 80 kd, when bound and chemically cross-linked to ¹²⁵I-TGF- β (Cheifetz et al., J. Biol. Chem. 265:20533-20538 (1990); Cheifetz and Massagué, J. Biol. Chem. 266:20767-20772 (1991); Massagué, J., Cell 5 69:1067-1070 (1992)). This receptor associates with the type I receptor (a 53 kd protein) to transduce the TGF- β signals; it is proposed that receptor I needs receptor II to bind TGF- β and that receptor II needs receptor I to mediate a signal (Laiho et al., J. Biol. Chem. 266:9108-10 9112 (1991); Laiho et al., J. Biol. Chem. 265:18518-18524 (1990); Wrana et al., Cell 71:1003-1014 (1992)). Transfection experiments with a recently cloned receptor I, also a serine/threonine kinase, further supports the view that receptor I must associate with receptor II to bind 15 TGF- β (Ebner et al., Science 260:1344-1348 (1993)).

The present invention provides a soluble endoglin-derived human polypeptide that binds TGF- β . The full-length soluble endoglin-derived polypeptide comprises a signal sequence that is cleaved during processing, the 20 561 amino acids of the extracellular domain of the mature endoglin polypeptide, an integral membrane protein, which consists of about 600 or 633 amino acids in total. Nucleic acid sequences encoding the human endoglin polypeptide are identified in Table 1 and Figure 13. The nucleic acid 25 sequences encoding the soluble endoglin-derived polypeptides are included within the sequences set forth in Table 1 (from about amino acid number 1 to about amino acid number 561) and Figure 13. Also provided by this invention is a vector having inserted therein the genomic DNA 30 molecule encoding endoglin. This vector was deposited under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure, with Coleccion Espanola de Cultivos Tipo (CECT), 46100 Burjasot 35 (Valencia), Spain on October 21, 1993, under CECT 4475. Accordingly, isolated genomic DNA encoding endoglin is

within the scope of this invention.

This invention also provides purified human endoglin polypeptides encoding two isoforms differing from each other in the cytoplasmic region. The "S-endoglin" has 14 amino acid residues in the cytoplasmic region and the "L-endoglin" has 47 amino acid residues in the cytoplasmic region. The 586 amino acids spanning the extracellular and transmembrane regions in the mature endoglin are identical.

As used herein, the term "purified" means that the molecule or compound is substantially free of contaminants normally associated with a native or natural environment. For example, the mature human proteins can be obtained from a number of methods. The methods available for the purification of membrane proteins include precipitation, gel filtration, ion-exchange, reverse-phase, and affinity chromatography. Other well-known methods are described in Deutscher et al., Guide to Protein Purification: Methods in Enzymology Vol. 182, (Academic Press 1990), which is incorporated herein by reference. Alternatively, a purified polypeptide of the present invention can also be obtained by well-known recombinant methods as described, for example, in Sambrook et al., Molecular Cloning: A Laboratory Manual 2d ed. (Cold Spring Harbor Laboratory 1989), also incorporated herein by reference. An example of this means for preparing soluble endoglin-derived polypeptide is to express nucleic acid encoding the soluble endoglin in a suitable host cell, such as a bacterial, yeast or mammalian cell, using methods well known in the art, and recovering the expressed soluble protein, again using methods well known in the art.

For the purpose of illustration, expression of soluble endoglin was achieved by excising a full length 2.3 kb endoglin cDNA fragment from pcEXV-L ENDO, Hind III linkers were added and, after digestion with Hind III, it

was subcloned into the multiple cloning site of pBluescript vector (Stratagene), followed by digestion with Tth 111I. (This enzyme cuts the coding region approximately 80 bp upstream of the transmembrane region.)

- 5 Synthetic, complementary oligomers were engineered to contain an in-frame stop codon and a Bam HI overhang, and were ligated to the linearized plasmid. After digestion with Hind III, and Bam HI, and purification of the 1.7 kb fragment, it was ligated to pCDNAI/Neo
10 (Invitrogen, San Diego, Figure 12). After cloning, the clones were screened by restriction enzyme analysis for the correct orientation and were found to be in the correct orientation. Verification of the cDNA insert was done using T7 and SP6 sequencing primers to confirm the presence
15 of the start and stop codons at the 5' and 3' ends of the insert..

5' End Primer

TAATACGACTCACTATAGGGAGACCCAAGCTTGGGGAATTCCGTGGACAGCATG

T7 PRIMER

HINDIII ECORI

Initiation

20 3' End Primer

AAGACCGTCTAGACGGATCCACTAG.....CTATAGTGTCACCTAAATG

BAMHI

SP6 PRIMER

- To express the construct, CHO-k1 cells (ATCC) were transfected with pCDNA/Neo to determine optimal
25 conditions required to generate stable transfectants resistant to G418. Five (5) x 10⁶ cells electroplated with 5 µg of DNA at a voltage of 300 volts and a time constant of 17.9 msec., using the 960 µF capacitor (Biorad Gene Pulse) yielded the most number of stable transfectants with
30 minimal cell death.

The following flow chart illustrates a means to express nucleic acid encoding soluble endoglin.

16

SOLUBLE ENDOGLIN

plasmid pcEXV-L ENDO
Eco R1 digest

5 Isolate 2.3 kb fragment

Add Hind III linker & Hind III digest

10 Subclone into pBluescript
MCS-Hind III
& selection on X-gal

15 pBluescript-L Endo

20 Digest with Tth 111I to linearize

Add complementary oligomers creating
stop codon & Bam HI overhang

25 TCTAGACG
AGATCTGCCTAG

30 Hind III Digest

Isolate 1.7 kb Hind III-BamHI fragment

35 Ligate with dephosphorylated
Hind III-Bam HI fragment of vector
pcDNAI/Neo

40 Transform E. coli MC1061/P3
selection on Tetracyclin & Ampicillin

The soluble polypeptide and biologically active fragments thereof can also be produced by chemical synthesis. Synthetic polypeptides can be produced using Applied Biosystems, Inc. Model 430A or 431A automatic polypeptide synthesizer and chemistry provided by the manufacturer. The soluble polypeptide can also be isolated directly from cells which have been transformed with the expression vectors described below in more detail.

As used herein, endoglin-derived polypeptide means a human polypeptide having the amino acid sequence substantially the same as the 633 amino acid sequence shown in Table 1, or the 658 amino acid sequence shown in Figure 13, or an active fragment thereof. As used herein the term "soluble endoglin-derived polypeptide" refers to a soluble, biologically active fragment of the human endoglin polypeptide expressed by the extracellular domain of the nucleic acid. As used herein, an "active fragment" or "biologically-active fragment" refers to any portion of an endoglin polypeptide that binds to TGF- β . Methods of determining whether a polypeptide can bind TGF- β are well known to those of skill in the art, for example, as set forth herein.

The invention also encompasses nucleic acid molecules which differ from that of the nucleic acid molecule shown in Table 1, e.g., the sequence shown in Figure 13, but which produce the same phenotypic effect. These altered, but phenotypically equivalent nucleic acid molecules are referred to as "equivalent nucleic acids." This invention also encompasses nucleic acid molecules characterized by changes in non-coding regions that do not alter the phenotype of the polypeptide produced therefrom when compared to the nucleic acid molecule described above. This invention further encompasses nucleic acid molecules which hybridize to the nucleic acid molecule of the subject invention or its complement. As used herein, the term

"nucleic acid" encompasses mRNA and cRNA as well as single and double-stranded genomic DNA, DNA and cDNA. In addition, as used herein, the term "polypeptide" encompasses any naturally occurring allelic variant thereof, such as S-endoglin and L-endoglin, as well as man-made recombinant forms.

This invention provides an isolated nucleic acid molecule encoding a soluble endoglin-derived polypeptide. As used herein, the term "isolated nucleic acid molecule" means a nucleic acid molecule that is in a form that does not occur in nature. One means of isolating a human endoglin nucleic acid is to probe a human cDNA expression library with a natural or artificially designed antibody to endoglin, using methods well known in the art (see Gougos, A. et al., J. Biol Chem. 265:8361 (1990)) and the Examples set forth below. DNA and cDNA molecules which encode human endoglin polypeptides can be used to obtain complementary genomic DNA, cDNA or RNA from human, mammalian or other animal sources. Isolated genomic DNA also is encompassed by this invention as described above. It is isolated by using the nucleic acid sequences of this invention and methods well known to those of skill in the art as described in Sambrook et al., supra.

The invention further provides an isolated nucleic acid molecule operatively linked to a promoter of RNA transcription, as well as other regulatory sequences. As used herein, the term "operatively linked" means positioned in such a manner that the promoter will direct the transcription of RNA off of the nucleic acid molecule. Examples of such promoters are SP6, T4 and T7. Vectors which contain both a promoter and a cloning site into which an inserted piece of DNA is operatively linked to that promoter are well known in the art. Preferable, these vectors are capable of transcribing RNA in vitro or in vivo. Examples of such vectors are the pGEM series

(Promega Biotec, Madison, WI).

This invention provides an expression or replication vector comprising this isolated nucleic acid molecule such as DNA, cDNA or RNA encoding a soluble endoglin-derived polypeptide. Examples of vectors are viruses, such as bacteriophages, baculoviruses and retroviruses, cosmids, plasmids (such as pcEXV-2) and other recombination vectors. Nucleic acid molecules are inserted into vector genomes by methods well known in the art. For example, insert and vector DNA can both be exposed to a restriction enzyme to create complementary ends on both molecules that base pair with each other and which are then joined together with a ligase. Alternatively, synthetic nucleic acid linkers can be ligated to the insert DNA that correspond to a restriction site in the vector DNA, which is then digested with a restriction enzyme that recognizes a particular nucleotide sequence. Additionally, an oligonucleotide containing a termination codon and an appropriate restriction site can be ligated for insertion into a vector containing, for example, some or all of the following: a selectable marker gene, such as neomycin gene for selection of stable or transient transfectants in mammalian cells; enhancer/promoter sequences from the immediate early gene of human CMV for high levels of transcription; transcription termination and RNA processing signals from SV40 for mRNA stability; SV40 polyoma origins of replication and ColE1 for proper episomal replication; versatile multiple cloning sites; and T7 and SP6 RNA promoters for in vitro transcription of sense and anti-sense RNA. Other means are available.

Also provided are vectors comprising a DNA molecule encoding an endoglin-derived polypeptide, or soluble fragment thereof, adapted for expression in a bacterial cell, a yeast cell, a mammalian cell and other animal cells. The vectors additionally comprise the

regulatory elements necessary for expression of the DNA in the bacterial, yeast, mammalian or animal cells so located relative to the DNA encoding soluble endoglin polypeptide as to permit expression thereof. Regulatory elements
5 required for expression include promoter sequences to bind RNA polymerase and transcription initiation sequences for ribosome binding. For example, a bacterial expression vector includes a promoter such as the lac promoter and for transcription initiation the Shine-Dalgarno sequence and
10 the start codon AUG (Sambrook et al. supra. 1989). Similarly, a eucaryotic expression vector includes a heterologous or homologous promoter for RNA polymerase II, a downstream polyadenylation signal, the start codon AUG, and a termination codon for detachment of the ribosome.
15 Such vectors can be obtained commercially or assembled by the sequences described in methods well known in the art, for example the methods described above for constructing vectors in general. Expression vectors are useful to produce cells that express the polypeptide.

20 This invention provides a mammalian cell containing a cDNA molecule encoding an endoglin-derived polypeptide or a soluble fragment thereof. An example is a mammalian cell comprising a plasmid adapted for expression in a mammalian cell. The plasmid has a cDNA
25 molecule encoding an endoglin-derived polypeptide and the regulatory elements necessary for expression of the polypeptide. Various mammalian cells may be utilized as hosts, including, for example, mouse fibroblast cell NIH3T3, CHO cells, HeLa cells, Ltk- cells, etc. Expression
30 plasmids such as those described supra can be used to transfect mammalian cells by methods well known in the art such as calcium phosphate precipitation, DEAE-dextran, electroporation or microinjection.

This invention provides a pharmaceutical
35 composition containing a pharmaceutical carrier and any of

a purified polypeptide, a purified soluble polypeptide, an active fragment thereof, or a purified, mature protein and active fragments thereof, alone or in combination with each other. These polypeptides or proteins can be recombinantly
5 derived, chemically synthesized or purified from native sources. As used herein, the term "pharmaceutically-acceptable carrier" encompasses any of the standard pharmaceutical carriers, such as a phosphate buffered saline solution, water and emulsions such as an oil/water
10 or water/oil emulsion, and various types of wetting agents. Any of these pharmaceutical compositions are useful in the methods described below or for the preparation of a medicament for treating the conditions described below.

Also provided are antibodies having specific
15 reactivity with the endoglin-derived TGF- β -binding polypeptides of the subject invention, such as anti-endoglin antibody 44G4 (Quackenbush, E.J., and Letarte, M.J., J. Immunol. 134:1276-1285 (1985)) or any antibody having specific reactivity to a TGF- β -binding endoglin
20 polypeptide. Active fragments of antibodies are encompassed within the definition of "antibody." The antibodies and fragments of the invention can be produced by any method known in the art. For example, polyclonal and monoclonal antibodies can be produced by methods well
25 known in the art, as described, for example, in Harlow and Lane, Antibodies: A Laboratory Manual (Cold Spring Harbor Laboratory 1988), which is incorporated herein by reference. The polypeptide can be used as the immunogen in generating such antibodies. Altered antibodies, such as
30 chimeric, humanized, CDR-grafted or bifunctional antibodies can also be produced by methods well known to those skilled in the art. Such antibodies can also be produced by hybridoma, chemical synthesis or recombinant methods described, for example, in Sambrook et al., supra. The
35 antibodies can be used for determining the presence or purification of the endoglin-derived polypeptide or soluble

fragment thereof, of the present invention. With respect to the detecting of such polypeptides, the antibodies can be used for in vitro diagnostic methods to determine the presence of endoglin or in vivo imaging methods.

5 Immunological procedures useful for in vitro detection of the target soluble endoglin-derived polypeptide in a sample include immunoassays that employ a detectable antibody. Such immunoassays include, for example, ELISA, Pandex microfluorimetric assay,
10 agglutination assays, radioimmunoassays, flow cytometry, serum diagnostic assays and immunohistochemical staining procedures which are well known in the art. An antibody can be made detectable by various means well known in the art. For example, a detectable marker can be directly or
15 indirectly attached to the antibody. Useful markers include, for example, radionuclides, enzymes, fluorogens, chromogens and chemiluminescent labels.

This invention provides a method of modifying a biological function mediated by the regulatory activity of
20 TGF- β which comprises contacting a suitable sample containing TGF- β with an effective amount of a biologically active endoglin-derived polypeptide, for example soluble endoglin, or a pharmaceutical composition described above.

As used herein, "an effective amount" refers to
25 an amount of the polypeptide sufficient to bind to TGF- β and thereby prevent or inhibit its regulatory activity. This method is especially useful for modifying the regulatory activity of TGF- β 1 or TGF- β 3. Examples of regulatory activities include, but are not limited to
30 stimulation of cell proliferation, cell growth inhibition, promotion of extracellular matrix proteins, and regulation of immune functions. TGF- β is known to be a potent chemoattractant for monocytes and can induce IL-1, TNF- α , TGF- β and surface Fc γ RIII, all of which are involved in the

inflammatory response. Conversely, TGF- β can deactivate macrophages by inhibiting the antimicrobial activity and the superoxide anion generation, and induce suppression of class II-restricted Ag presentation by macrophages.

- 5 The method can be practiced in vitro or in vivo.
If the method is practiced in vitro, contacting is effected by incubating the sample with a polypeptide, a protein or a pharmaceutical composition as described above.

- 10 In vitro the novel nucleic acid molecules and antibodies of this invention are useful to detect and quantify the amount of TGF- β in a sample isolated from a subject, such as a human patient. The detection of TGF- β is useful to monitor the progression of a disease related to overexpression of TGF- β , e.g., glomerulonephritis.

- 15 However, in a preferred embodiment the contacting is effected in vivo by administering a polypeptide, a protein or a pharmaceutical composition, as described above, to a subject, e.g., a human patient.

- 20 Methods of administration are well known to those of skill in the art and include, but are not limited to administration orally, intravenously or parenterally. Administration will be in such a dosage such that the regulatory activity is effectively modified. Administration can be effected continuously or
25 intermittently such that this amount is effective for its intended purpose.

- 30 This invention also provides a method of treating a pathologic condition caused by a TGF- β -regulated activity comprising contacting the TGF- β with any of a purified soluble endoglin-derived polypeptide, an active fragment thereof, an endoglin-derived polypeptide or an active fragment thereof. The TGF- β is bound with said polypeptide

to thereby treat the pathologic condition mediated by TGF- β regulatory activity. As used herein, "pathologic conditions" refers to any pathology arising from TGF- β -induced regulatory activity, for example, inflammation, 5 rheumatoid arthritis, inflamed skin lesions, scar tissue formation, lung fibrosis, liver fibrosis, atherosclerosis, and glomerulonephritis. Growth and proliferation of mesenchymal cells is stimulated by TGF- β , however some tumor cells may also be stimulated thus using TGF- β as an 10 autocrine growth factor. An example of inhibitory conditions are the prevention of new cell growth to assist in repair of tissue damage, for ulceration and immunosuppression. The stimulation of extracellular matrix production by TGF- β is essential for wound healing. 15 However, in some cases, the TGF- β response is uncontrolled and an excessive accumulation of extracellular matrix results. An example of excessive accumulation of extracellular matrix is glomerulonephritis. An additional example of a pathology is cancer.

20 In one embodiment, the method is practiced by administering to a subject, e.g., a human patient or a mammal, an effective amount of a purified endoglin protein or an endoglin-derived soluble polypeptide or a biologically active fragment thereof, or the pharmaceutical 25 composition described above. Methods of administration are outlined supra.

Also provided by this invention is a method of inhibiting the activity of endoglin by contacting endoglin with an effective amount of a polypeptide capable of 30 binding to endoglin to bind endoglin, thereby inhibiting the activity of endoglin. As used herein, the term "a polypeptide capable of binding to endoglin" means any substance capable of forming a complex with endoglin, for example, TGF- β 1 or TGF- β 3, or an active fragment thereof. 35 An active fragment is an amino acid sequence corresponding

to a fragment of TGF- β 1 or TGF- β 3 that retains the ability to bind endoglin. Methods of making such fragments are well known to those of skill in the art as are methods of determining the binding activity of the fragments. Also encompassed by this invention are polypeptides that retain their activity to bind to endoglin, but no longer mediate the biological response corresponding to the binding of a functional ligand to the receptor is destroyed. Thus, these "mutated" polypeptides can act as antagonists to the biological function mediated by the ligand to endoglin by blocking the binding of normal, functioning ligands to endoglin on the cell.

This invention also encompasses the use of the compositions defined above for the preparation of medicaments to modify a biological function regulated by TGF- β . These biological functions are described above in detail.

It is understood that modifications which do not substantially affect the activity of the various molecules of this invention are also included within the definition of said molecules.

The following examples are intended to illustrate but not limit the present invention.

**A. ISOLATION OF HUMAN ENDOGLIN PROTEIN
AND NUCLEIC ACID ENCODING SAME**

EXAMPLE I

Cell Culture and Transfections

Human umbilical vein endothelial cells (HUVEC, CRL 1730, ATCC) were maintained in α -minimal essential media supplemented according to supplier's instructions or prepared from umbilical veins as previously described

(Gougos, A. et al., J. Immunol. 141:1925 (1988)). Similar results were obtained using cells from either source. COS-M6 cells, maintained in Dulbecco's modified Eagle's medium supplemented with 10% bovine serum, were transfected with
5 a cDNA encoding full-length endoglin ligated into the EcoRI site of the mammalian expression vector pcEXV (Miller, J. et al., J. Exp. Med. 164:1478 (1986)) or with a control vector without cDNA insert (pcMV5; Lopez-Casillas, F. et al., Cell 67:785 (1991)) by the DEAE-dextran-chloroquine
10 procedure (Seed, B., et al., P.N.A.S. USA 84:3365 (1987)). Twenty-four (24) hours post-transfection, cells were trypsinized and reseeded into multicluster dishes and allowed to grow an additional 48 hours before being affinity-labeled with ^{125}I -TGF- β 1 as described below.

15

EXAMPLE II

Receptor Affinity Labeling and Immunoprecipitation

TGF- β 1 and TGF- β 2 were purchased from R & D Systems (Minneapolis, MN) and TGF- β 3 was obtained from Oncogene Science (Manhasset, NY). ^{125}I -TGF- β 1 used in these
20 studies was prepared by the chloramine-T method as previously described (Cheifetz, S. et al., J. Biol. Chem. 265:20533 (1990)) or purchased from Amersham Corp.; both preparations gave identical results. The conditions for affinity labeling cell monolayers with ^{125}I -TGF- β 1 and
25 disuccinimidyl suberate (Pierce Chemical Co.) have been described previously (Massague, J., Methods Enzymol. 146:174 (1987)). The concentrations of ^{125}I -TGF- β 1 and competing unlabeled ligands used for each experiment are indicted in the figure legends. Triton X-100 extracts of
30 the affinity-labeled cells were either analyzed directly on sodium dodecyl sulfate-polyacrylamide gels (SDS-PAGE) or first incubated with monoclonal antibody (mAb) 44G4 directed against human endoglin (Quackenbush, E.J. et al., J. Immunol. 134:1276 (1985)) or with control antibody (see
35 below). For immunoprecipitations, detergent extracts were

diluted with an equal volume of phosphate-buffered saline containing 1% Triton X-100 and precleared by incubation for 20 min at 4°C with protein G-Sepharose (Pharmacia LKB Biotechnology Inc.) prior to overnight incubation at 4° C. with mAb 44G4. Immune complexes were collected by incubation with protein G-Sepharose for 1 hour at 4° C. For some experiments, mAb 44G4 was used coupled to Sepharose. The immunoprecipitates were washed three times (saline with 1% Triton X-100) and then resolved by SDS-PAGE in the presence or absence of dithiothreitol (DTT) and visualized by autoradiography. Irrelevant mAb (44D7) used in control experiments to monitor specificity of the immunoprecipitations did not immunoprecipitate any affinity-labeled bands.

15

EXAMPLE III

SDS-PAGE and 2D-Gel Analysis

Analysis of the affinity-labeled profile of HUVEC revealed that, like vascular endothelial cells from other sources, these cells have little or no betaglycan, which characteristically migrates as a diffuse band between 200 and 400 KDa on reducing SDS-PAGE (Figure 2A). Instead, HUVEC expressed a disulfide-linked cell surface protein that, together with TGF- β receptors I and II, was affinity-labeled by crosslinking with 125 I-TGF- β 1. Receptors I and II were detected in HUVEC as labeled complexes of approximately 65 KDa and 100 KDa, respectively, which is similar to the size of these labeled receptors reported for other human cell lines. Comparison of the relative migration of the affinity-labeled proteins fractionated on SDS-PAGE revealed that the major affinity-labeled proteins of HUVEC migrated between 95-120 KDa on reducing gels whereas on non-reducing gels the major affinity-labeled proteins migrated between 100-110 KDa (presumed to be receptor II) and at 180 KDa and above (endoglin) (Figure 2). This pattern indicated the presence of disulfide-

linked TGF- β -binding proteins.

Resolution of these disulfide-linked TGF- β 1 binding proteins on two-dimensional gels (Figure 2B) confirmed that the disulfide-linked complexes (probably dimers and higher order oligomers) contained subunits of approximately 95 KDa (value estimated by subtracting the cross-linked TGF- β 1 monomer mass 12.5 KDa from the reduced 110 KDa affinity-labeled complex). Together with the type II receptor, the disulfide-linked TGF- β 1-binding proteins are the major affinity-labeled species expressed by HUVEC.

EXAMPLE IV

Immunoprecipitation with anti-endoglin mAb

To determine whether the disulfide-linked TGF- β -binding protein on endothelial cells was endoglin, affinity-labeled HUVEC extracts were immunoprecipitated with monoclonal antibody (mAb) 44G4 which is specific for human endoglin (Georgi, L.L. et al., Cell 61:635 (1990); MacKay, K. et al., J. Biol. Chem. 266:9907 (1992); Merwin, J.R. et al., Am. J. Pathol. 138:37 (1991)). Electrophoretic analysis of these immunoprecipitates revealed a labeled protein complex whose subunit structure was similar to that of endoglin (Figure 3A). Thus, under reducing conditions, a major affinity-labeled band of approximately 110 KDa was seen which migrated as complexes of 180 KDa and greater than 200 KDa when analyzed under non-reducing conditions. The higher order oligomers might contain multiple endoglin molecules crosslinked by TGF- β 1, itself a disulfide-linked dimer. Repeated immunoprecipitation with 44G4-IgG-Sepharose completely depleted these labeled species from cell extracts (Figure 3B). No affinity-labeled bands were immunoprecipitated from three other human cell lines (A549, Hep G2, MCF-7), which lack endoglin and were used as negative-controls for these experiments. Monoclonal antibodies specific to human

endoglin thus demonstrate that endoglin is a major TGF- β -binding protein in human vascular endothelial cells.

EXAMPLE V

Ectopic Expression of Endoglin in Cells

5 The identity of this dimeric TGF- β -binding protein of HUVEC with endoglin was confirmed by ectopically expressing the full-length endoglin cDNA in COS monkey kidney cells. After affinity-labeling with ^{125}I -TGF- β 1, a labeled species with the characteristics of endoglin could
10 be specifically precipitated by mAb 44G4 only from the detergent extracts of endoglin transfectants (Figure 4). Differences in glycosylation likely account for the smaller size of endoglin expressed in COS cells relative to endogenous endoglin of HUVEC.

15 B. ISOLATION OF S- AND L-ENDOGLIN

EXAMPLE VI

cDNA Cloning and Sequencing of Endoglin

Approximately 2.5×10^5 clones from a λ gt10 cDNA library prepared from PMA-treated myelomonocytic human cell
20 line HL60 (Corbi, A.L. et al., EMBO J. 6:4023 (1987)). (50,000 pfu/150 mm dish) were screened with 700 bp PstI fragment from endoglin cDNA (Gougos, A. and Letarte, M.J., J. Biol. Chem. 265:8361 (1990)). Thirteen hybridizing clones were isolated after three rounds of plaque
25 purification with sizes ranging from 2.4-3.1 kb. The longest clone (clone 3.3; 3073 bp) was subcloned into pUC13 and sequenced using the dideoxy chain termination method (Sanger, F. et al., PNAS USA 74:5463 (1977)).

Clone 3.3 in pUC13 was digested with BbrPI and
30 BamHI. Endoglin fragment was made blunt and inserted into the mammalian expression vector pcEXV (Miller, J. and

Antibodies, Immunofluorescence, Immunoprecipitations and Immunoblotting

Endoglin-specific monoclonal antibodies used were 8E11 (Lastres, P. et al., Eur. J. Immunol. 22:393 (1992)) of the IgM class and 44G4 (Gougos, A. and Letartè, M., J. Immunol. 141:1925 (1988)) of the IgG1 subclass. Control monoclonal antibodies were HC1/1 (anti-CD11c) of the IgG1 subclass and X63. FCM and immunoprecipitation analyses were performed as described in Lastres, P. et al., Eur. J. Immunol. 22:393 (1992), incorporated herein by reference. For immunoblotting studies, 5×10^6 cells were lysed in 250 μ l of 1% Triton X-100, 1 mM PMSF in PBS for 30 minutes at 4°C. Insoluble material was removed by centrifugation at 100,000 x g for 1 hour in a Beckman TL100 centrifuge. Proteins contained in the supernatant were separated by SDS-PAGE under nonreducing conditions using a minigel system and transferred to nitrocellulose membranes (Hybond, Amersham). Membranes were incubated first with a blocking solution (10% fetal calf serum, 0.5% Tween-20, 1 M glucose and 10% glycerol in PBS), followed by incubation with the monoclonal antibody 44G4. The presence of endoglin was revealed using a chemiluminescence assay (ECL detection

kit, Amersham).

EXAMPLE VIII

Reverse Transcription-Polymerase Chain Reaction (RT-PCR)

Total cellular RNA was isolated using guanidinium isothiocyanate and a cesium chloride ultracentrifugation step (Chirgwin, J.M. et al., Biochemistry 18:5294 (1979)). Poly (A)+ RNA was purified by oligo (dT) affinity chromatography (Maniatis, T. et al., "Molecular Cloning: A Laboratory Manual" Cold Spring Harbor Laboratory, Cold Spring Harbor (1982). Single-stranded cDNA was synthesized from poly (A+) RNA (0.5 µg/40 µl reaction), using AMV reverse transcriptase. Five microliters of cDNA was used for a 50 µl PCR reaction (Gilliland, G. et al., In: [Innis, M.A., Gelfand, D.H., Sninsky, J.J. and White, T.J. (Eds.)) "PCR Protocols" Academic Press, San Diego 1990, p. 60.). Oligonucleotide primers used for amplification were E#12 (nucleotides 1945-1964), E#11R (reverse complement of nucleotides 2475-2494), E#14 (nucleotides 2136-2158), and E#15 (reverse complement of nucleotides 2247-2273). Amplifications were performed in 1 X Taq buffer (Promega) with 0.2 mM each dATP, dCTP, dGTP, dTTP at 0.25 µM concentration of each primer and 0.25 U/50 µl of Taq DNA polymerase (Promega). Amplification was carried out in a thermal cycler as follows: 5 minutes at 95°C; 35 cycles of 45 seconds at 94°C, 45 seconds at 54°C, and 1 minute at 72°C for oligonucleotide pair E#12-E#11R or 35 cycles of 45 seconds at 94°C, 45 seconds at 68°C and 1 minute at 72°C for pair E#14-E#15 and then 10 minutes at 72°C. Control reactions, identical except for the omission of reverse transcriptase, were performed simultaneously. Amplification products were analyzed by agarose gel electrophoresis and ethidium bromide staining.

EXAMPLE IX

Receptor Affinity Labeling

TGF- β 1 was purchased from R&D Systems (Minneapolis, MN) and 125 I-TGF- β 1 was obtained from Amersham (Oakville, Canada) at a specific activity of 2000 Ci/mmol. Parental mouse L cells, transfectant mouse L cells expressing S-endoglin or L-endoglin, and mock transfectant L cells were grown at confluence (5×10^6 cells per plate) and incubated with 100 pM of 125 I-TGF- β 1 for 4 hours with and without 4 nM of competing unlabeled TGF- β 1. Cells were washed and cross-linked at a final concentration of 0.16 mM disuccinimidyl suberate (DSS) (Pierce Chemical Co.) in a buffer containing 128 mM NaCl, 5 mM KCl, 5 mM MgSO₄, 1.2 mM CaCl₂, 50 mM Hepes, pH 7.5 for 15 minutes at 4°C. Cells were washed four times and solubilized directly on the petri dishes with a minimum of solubilization buffer containing 1% Triton X-100 and proteolytic inhibitors as described previously (Massagué, J., Methods Enzymol. 146:174 (1987)). The extracts were subjected to immunoprecipitation followed by SDS-PAGE analysis.

RESULTS

Isolation and Characterization of Endoglin cDNA Clones:
Identification of a Cytoplasmic Variant

Full-length cDNA clones were derived from a λ gt10 library prepared from PMA-treated HL60 cells. Clone number 3.3 was selected for further characterization due to the large size of the insert. The partial sequence of the 3073 bp cDNA insert from clone 3.3 is depicted in Figure 7. This clone contains 281 bp of 5' untranslated region followed by an open reading frame of 1875 bp. The predicted protein sequence shows that the endoglin leader peptide contains 25 amino acids (aa), followed by 561 residues at the extracellular portion and a transmembrane

region spanning 25 amino acids as expected (Gougos, A. and Letarte, M.J., J. Biol. Chem. 265:8361 (1990)). However, clone 3.3 contains a 135 bp segment inserted within the previously known cDNA sequence, starting at nucleotide (nt) 2134 (Figure 8). The first 21 nucleotides of the insert are in frame with the preceding sequence and code for a new sequence of 7 amino acids. This sequence replaces the C-terminal 40 amino acids of endoglin, thus leading to a new cytoplasmic tail of 14 amino acids different from the 47 residues previously reported (Gougos, A. and Letarte, M.J., J. Biol. Chem. 265:8361 (1990)) (Figure 8), suggesting the existence of two alternative forms of endoglin. The predominant form of endoglin in the myelomonocytic cell line HL60 seems to be the one containing 47 residues in the cytoplasmic domain since this sequence was present in 12 out of 13 clones analyzed. The isoforms with the 14 and 47 amino acid cytoplasmic tails, and the corresponding cDNAs, are referred to as S-endoglin and L-endoglin, respectively.

Expression of two different forms of endoglin

To characterize further the two alternative forms, independent constructs corresponding to the short and long forms of endoglin were inserted into the expression vector pcEXV and transfected into mouse fibroblasts (Figure 9). Both, S-endoglin and L-endoglin were highly expressed on the cell surface as determined by FCM (Figure 9). In addition, metabolic labeling of the transfectants followed by immunoprecipitation, revealed a 170-kDa (L-endoglin) or a 160-kDa (S-endoglin) protein specifically recognized by anti-endoglin monoclonal antibody (Figure 9). The distinct size of S- and L-endoglin in the transfectants was also detected by immunoblotting analysis under non-reducing conditions (Figure 9). In these experiments, L-endoglin showed the same M_r as the endoglin detected on U937 cells, suggesting that this is the predominant form in the promonocytic cell

line.

Cell surface radiolabeling of the transfectants followed by immunoprecipitation demonstrated that both isoforms are expressed as disulfide linked homodimers (Figure 9), indicating that cysteine residue present in the extracellular portion are mediating the interchain disulfide bond.

Differential expression
of S-endoglin and L-endoglin mRNA

10 The individual expression of S- and L-endoglin mRNA was analyzed by RT-PCR. Amplification with two primers derived from the unique 3' region of S-endoglin cDNA generated the expected 137 bp product on S-endoglin transfectants and on PMA-treated monocytic lines, 15 endothelial cells and placenta (Figure 10), demonstrating the existence of this isoform in several cell types. When primers common to L-endoglin and S-endoglin were used, the L-endoglin specific fragment of 411 bp could be amplified on PMA-treated monocytic cell lines, placenta and 20 endothelial cells, whereas the S-endoglin fragment of 546 bp was only amplified on control S-endoglin transfectants (Figure 10). When competitive templates are amplified by PCR, an abundant template can suppress the amplification of a less abundant one (Gilliland, G. et al., In: [Innis, 25 M.A., Gelfand, D.H., Sninsky, J.J. and White, T.J. (Eds.) "PCR Protocols", Academic Press, San Diego 1990, p. 60]. Therefore, these results indicate that L-endoglin is the predominant form and S-endoglin is expressed at lower levels in these cell types. This is in agreement with the 30 presence in PMA-treated U937 cells of an endoglin molecule of M_r similar to that observed on the transfectant L-endoglin (Figure 9), assuming similar glycosylation levels on both cell types.

L-endoglin and S-endoglin bind to TGF- β 1

Since endoglin has been found to be a component of the receptor system for TGF- β , Cheifetz, S. et al., J. Biol. Chem. 267:19027 (1992), it was of interest to analyze the ability of each isoform to bind this ligand. Transfectants expressing short and long forms of endoglin were compared to the parental L cells and to the mock transfectants for their ability to bind TGF- β 1. Immunoprecipitation analysis allowed the identification of the TGF- β 1-endoglin complex (Figure 11). Under non-reducing conditions, the dimers of endoglin were seen as radiolabelled bands of M_r 170 kDa (short form) and 175 kDa (long form). An oligomer of endoglin is seen in both cases migrating with a M_r > 270 kDa. A similar complex was also observed previously in cross-linking experiments with human endothelial cells and might represent endoglin molecules cross-linked by TGF- β 1, itself a dimer (Roberts, A.B. and Sporn, M.B. In: Sporn, M.B. and Roberts, A.B. (Eds.), "Peptide growth factors and their receptors" Springer-Verlag, Heidelberg 1990, p. 419). Under reducing conditions, major bands of 97 kDa and 107 kDa were seen with the short form of endoglin and bands of 102 kDa and 112 kDa were observed with the long form. These doublets have been observed previously for endothelial cells and might represent endoglin bound to the monomer (12.5 kDa) or dimer (25 kDa) of TGF- β 1 (Roberts, A.B. and Sporn, M.B. In: Sporn, M.B. and Roberts, A.B. (Eds.), "Peptide growth factors and their receptors" Springer-Verlag, Heidelberg 1990, p. 419.; Cheifetz, S. and Massagué, J., J. Biol. Chem. 266:20767 (1991)). By subtracting the contribution of TGF- β 1, one can estimate a molecular weight of 85 kDa for S-endoglin and 90 kDa for L-endoglin.

DISCUSSION

Sequence analysis demonstrated the existence of two different cDNA variants named L-endoglin and S-endoglin. These two isoforms are coexpressed by myeloid cells, endothelium and placenta, although the majority of the transcripts synthesized apparently correspond to the L-endoglin isoform. The mechanism by which the two isoforms are generated remains to be determined. Most likely, both isoforms are generated by alternative splicing. In fact, consensus sequences of donor/acceptor sites (GT, AG) at the 5' and 3' ends and branch point of the lariat (CTGAC), have been found on the novel 135 bp insert of S-endoglin cDNA (Figure 7). A similar example of cytoplasmic variants generated by a "retained intron" splicing mechanism, have been recently reported for the activin receptor of the TGF- β receptor family (Attisano, L. et al., Cell 68:97 (1992)).

The behavior of the two endoglin isoforms was analyzed by transfection studies. Both forms are expressed on the cell surface as disulfide linked homodimers, indicating that the cysteine residues present in the extracellular region are responsible for the dimerization. In spite of the different cytoplasmic domains, both forms behave as TGF- β binding proteins.

S-endoglin, L-endoglin and betaglycan contain 22 identical residues in the transmembrane domain and the adjacent cytoplasmic region (Figure 9). This delineates a first conserved motif between betaglycan and the two forms of endoglin, which contains two tyrosine residues and two Ser/Thr residues whose phosphorylation status remain to be analyzed. A second region of high identity in the cytoplasmic tail is shared only by the long form of endoglin and betaglycan. The high content (40%) of Ser/Thr residues in this motif, and the absence of Tyr suggests that this region might undergo phosphorylation by a Ser/Thr

kinase. Interestingly, the cytoplasmic domain of the TGF- β receptor II, displays Ser/Thr kinase activity (Lin, H.Y. et al., Cell 68:775 (1992)).

Although the invention has been described with
5 reference to the disclosed embodiments, it should be understood that various modifications can be made without departing from the spirit of the invention. Accordingly, the invention is limited only by the following claims.

What is claimed is:

1. An isolated nucleic acid molecule encoding an endoglin polypeptide or a biologically active fragment thereof.
- 5 2. A nucleic acid molecule encoding a soluble endoglin-derived polypeptide or a biologically active fragment thereof.
3. The nucleic acid of claim 1 or 2, wherein the nucleic acid encodes a human endoglin polypeptide.
- 10 4. The nucleic acid molecule of claim 1 or 2, wherein the nucleic acid is genomic DNA, cDNA, mRNA or cRNA.
5. The nucleic acid molecule of claim 4, wherein the genomic DNA is designated CECT 4475.
- 15 6. The nucleic acid molecule of claim 1 or 2 operatively linked to a promoter of RNA transcription.
7. A vector containing the nucleic acid molecule of claim 5.
8. A host cell containing the vector of claim
- 20 7.
9. The host cell of claim 8, wherein the cell is a procaryotic cell or a eucaryotic cell.

10. A method of preparing an endoglin-derived polypeptide or active fragment thereof, comprising:

- a. inserting a nucleic acid molecule encoding an endoglin-derived polypeptide or active fragment thereof
5 into a suitable expression vector;
 - b. inserting the resulting vector into a suitable host cell;
 - c. inducing the resulting host cell to express the endoglin-derived polypeptide or active fragment
10 thereof; and
 - d. purifying the resulting endoglin-derived polypeptide so produced.
11. Soluble endoglin-derived polypeptide.
12. Endoglin-derived polypeptide or a
15 biologically active fragment thereof produced by the method of claim 10.
13. A pharmaceutical composition comprising the polypeptide of claim 11 or 12 and a pharmaceutically acceptable carrier.
- 20 14. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a purified and isolated human endoglin-derived polypeptide or a biologically active fragment thereof substantially free of other host cell proteins.

15. A method of modifying a biological function mediated by the regulatory activity of TGF- β which comprises contacting TGF- β with an effective amount of an endoglin-derived polypeptide or a biologically active
5 fragment thereof, thereby modifying the biological function.

16. The method of claim 15, wherein TGF- β is TGF- β 1.

17. The method of claim 15, wherein TGF- β is
10 TGF- β 3.

18. The method of claim 15, wherein the contacting is effected in vitro.

19. The method of claim 15, wherein the contacting is effected in vivo.

20. The method of claim 15, wherein the regulatory activity is stimulation of cell proliferation or cell growth inhibition.

21. The method of claim 15, wherein the regulatory activity is promotion of extracellular matrix
20 production.

22. The method of claim 15, wherein the regulatory activity is regulation of immune function.

23. The method of claim 15, wherein the polypeptide is a purified human endoglin-derived
25 polypeptide comprising a disulfide-linked homodimer of about 80 to about 95 kDa subunits or a biologically active fragment thereof.

24. The method of claim 23, wherein the purified polypeptide is purified and isolated human endoglin-derived protein substantially free of other human proteins.

25. The method of claim 15, wherein the polypeptide is a soluble endoglin-derived polypeptide.

26. A method of treating a pathologic condition caused by TGF- β regulated cell growth stimulation which comprises administering to a subject an effective amount of the pharmaceutical composition of claim 13 or 14 to bind to the TGF- β thereby treating the pathologic condition caused by cell growth stimulation.

27. The method of claim 26, wherein TGF- β is TGF- β 1.

28. The method of claim 26, wherein TGF- β is TGF- β 3.

29. The method of claim 26, wherein the subject is a human patient.

30. A method of treating a pathologic condition caused by TGF- β regulated inhibition of cell growth which comprises administering to a subject an effective amount of the pharmaceutical composition of claim 13 or 14 to bind to the TGF- β , thereby treating the pathologic condition caused by inhibition of cell growth.

31. The method of claim 30, wherein TGF- β is TGF- β 1.

32. The method of claim 30, wherein TGF- β is TGF- β 3.

33. The method of claim 30, wherein the pathologic condition is ulceration or immune suppression.

34. The method of claim 30, wherein the subject is a human patient.

5 35. The method of treating a pathologic condition caused by TGF- β regulated promotion of extracellular matrix accumulation which comprises administering to a subject an effective amount of the pharmaceutical composition of claim 12 or 13 to bind TGF- β
10 thereby treating the pathologic condition caused by promotion of extracellular matrix.

36. The method of claim 35, wherein the subject is a human patient.

15 37. The method of claim 35, wherein the pathologic condition is inflammation, rheumatoid arthritis, inflamed skin lesions, scar tissue formation, lung fibrosis, liver fibrosis, atherosclerosis or glomerulonephritis.

20 38. A method of inhibiting the activity of endoglin which comprises contacting endoglin with an effective amount of a polypeptide capable of binding to endoglin to bind endoglin, thereby inhibiting the activity of endoglin.

25 39. The method of claim 38, wherein the polypeptide is a TGF- β .

40. The method of claim 38, wherein the polypeptide has an amino acid sequence corresponding to a fragment of a TGF- β having the ability to bind endoglin.

41. The method of claim 39, wherein the TGF- β is TGF- β 1 or TGF- β 3.

42. The method of claim 40, wherein the TGF- β is TGF- β 1 or TGF- β 3.

5 43. The method of claim 38, wherein the polypeptide is an anti-endoglin antibody.

44. The method of claim 43, wherein the antibody is a monoclonal antibody.

10 45. The method of claim 38, wherein the contacting is effected in vivo.

46. The method of claim 38, wherein the contacting is effected in vitro.

- 1 / 13

FIGURE 1






FIGURE 2



- 3 / 13

FIGURE 3



- 4 / 13

FIGURE 4






FIGURE 5



pcNeoSolEND 8.7 Kb



Vector pcDNA1/Neo
7.0 Kb

[illegible]

FIGURE 9







FIGURE 10




FIGURE 11





FIGURE 12



1 : TGGGCGCAGGACTGCTGCTGTCACTGCCATTCATTGAAGCCGACGACCCTCTCCCGCCCATCTTTCGGACAGCAACTCCAGCCAGCCC
91 : CCGCTCTCTGTGTCCACTTCTCTGACCCCTCGCCCCCACCCAGAAAGCTGGAGCAGGAGCGCTGCTTCGCGCCCTGCTCTCCCTT
181 : CGGCTCCCGCTGGAGCGCCAGCCGCGCCCGGTGCCCCCGCCGAGCCCTGCCATCGACACAGGATAAGGCCCCAGCAGCCGCCCCAAC
271 : GTGGACAGCATGGACCCTGCGCAGCTTCTCTTGCTTGGCTCTGCTGCTGCAGCTTCAGCCCCAAGCTCTGTGAAAAACA
.....MetAspArgGlyThrLeuProLeuAlaValAlaLeuLeuAlaSerCysSerLeuSerProThrSerLeuAlaGluThr

361 : GTCCATTGTGACCTTCAGCCTGTGGGCCCCGAGAGGGGGGAGGTGACATATACCACTAGCCAGGCTCGAAGGGCTGCTGSGCTCAGGCC
28 : ValHisCysAspLeuGlnProValGlyProGluArgGlyGluValThrTyrThrThrSerGlnValSerLysGlyCysValAlaGlnAla

451 : CCCATGCCATCTTGAAGTCCATGTCTCTTCTGGAGTTCCCAACGGGGCCGTCACAGCTGGAGTGACTCTCCAGGATCCAAGCAA
59 : ProAsnAlaIleLeuGluValHisValLeuPheLeuGluPheProThrGlyProSerGlnLeuGluLeuThrLeuGlnAlaSerLysGln

541 : AATGGCACTTGGCCCCGAGAGGTGCTTCTGTCTCAGTGTAAACAGCAGTGTTCTTCTGCATCTCCAGGCCCTGGGAATCCCCTGCAC
98 : AsnGlyThrTrpProArgGluValLeuLeuValLeuSerValAsnSerSerValPheLeuHisLeuGlnAlaGlyIleProLeuHis

631 : TTGGCCTACAATTCCAGCCTGGTCACCTTCCAAGAGCCCCGGGGTCAACACCACAGAGCTGCCATCTTCCCCAAGACCCAGATCTCTT
128 : LeuAlaTyrAsnSerSerLeuValThrPheGlnGluProProGlyValAsnThrThrGluLeuProSerPheProLysThrGlnIleLeu

721 : GAGTGGCGAGCTGAGAGGGGGCCCATCCTCTGCTGCTGAGCTGAATGACCCACAGCATCTCTCTCCGACTGGGCCAAGCCAGGGG
148 : GluTrpAlaAlaGluArgGlyProIleThrSerAlaAlaGluLeuAsnAspProGlnSerIleLeuLeuArgLeuGlyGlnAlaGlnGly

811 : TCACCTGCTCTTCTGCATGCTTGAAGCCAGCCAGGACATGGGGCCGACGCTCGAGTGGCGCCGCGTACTCCAGCTTGGTCCGGGGTGC
178 : SerLeuSerPheCysMetGluLeuAlaSerGlnAspMetGlyArgThrLeuGluTrpArgProArgThrProAlaLeuValArgGlyCys

901 : CACTTGGAAAGCGTGCCCGGCCACAAAGGAGCGGCACATCTGAGGCTCTCGCGGCCCATCGSCCGGGCCCCGGACGGTGAAGTGAAG
208 : HisLeuGluGlyValAlaGlyHisLysGluAlaHisIleLeuArgValLeuLeuGlyHisSerAlaGlyProArgThrValThrValLys

991 : GTGGAAC TGAGCTGGCGCCGCGGATCTCSATGCCGTCTCATCTCGAGGGTCCCCCTACGTTGCTCTGGCTCATCGACGCCAACCAC
238 : ValGluLeuSerCysAlaProGlyAspLeuAspAlaValLeuIleLeuGlnGlyProProTyrValSerTrpLeuIleAlaSerAlaAsnHis

1081 : AACATGCAGATCTGGACCCTGAGAGAATACTCTCTCAAGATCTTTCCAGAGAAAACATTCTGGCTTCAAGCTCCACGACACACCTCAA
268 : AsnMetGlnIleTrpThrThrGlyGluTyrSerPheLysIlePheProGluLysAsnIleArgGlyPheLysLeuProAspThrProGln

1171 : GGCTCTCTGGGGAGGCCCGGATGCTCAATGCCAGCATTGTGGCATCTCTCGTGGAGCTACCGCTGGCCAGCATTTGCTCACTTCATGCC
298 : GlyLeuLeuGlyGluAlaArgMetGluLeuAsnAlaSerIleValAlaSerPheValGluLeuProLeuAlaSerIleValSerLeuHisAla

1261 : TCCAGCTGCGGTGGTAGGCTCCAGACCTACCCCGCCAGCAGCAGCATCTCCCAAGGACACTTGTAGCCCGGAGCTGCTCATGTCC
328 : SerSerCysGlyGlyArgLeuGlnThrSerProAlaProIleGlnThrThrProLysAspThrCysSerProGluLeuMetSer

1351 : TTGATCCAGACAAAGTGTGCCGACGACGCCATGACCTGGTACTAAAGAAGAGCTTGTTCGCGATTTCGAAGTGCACCATCAGGGCGCTG
358 : LeuIleGlnThrLysCysAlaAspAspAlaMetThrLeuValLeuLysLysGluLeuValAlaHisLeuLysCysThrIleThrGlyLeu

1441 : ACCTTCTGGGACCCAGCTGTGAGGCGAGGACAGGGGTGACAAGTTTGTCTTGGCAGTGCTTACTCCAGCTGTGGCATGCAGGTGTCA
388 : ThrPheTrpAspProSerCysGluAlaGluAspArgGlyAspLysPheValLeuArgSerAlaTyrSerSerCysGlyMetGlnValSer

1531 : GCAAGTATGATCAGCAATGAGGCGGGTGGTCAATATCTGTCGAGCTCATCACACAGCGGAAAAAGGTGCACCTGCCCTCAACATGGACAGC
418 : AlaSerMetIleSerAsnGluAlaValValAlaIleLeuSerSerSerProGlnArgLysLysValHisCysLeuAsnMetAspSer

1621 : CTCTCTTTCAGCTGGGCTCTACTCTCAGCCACACTCTCTCCAGGCTCTCAACACCATCGAGCCGGGAGCAGAGCTTGTGCGAGGT
448 : LeuSerPheGlnLeuGlyLeuTyrLeuSerProHisPheLeuGlnAlaSerAsnThrIleGluProGlyGlnGlnSerPheValGlnVal

1711 : AGAGTGTCCCATCCGTTCTCGAGTTCTGCTCCAGTTAGACAGCTGCCACCTGGACTTGGGGCTGAGGAGGCCAGCTGGAACCTCATC
478 : ArgValSerProSerValSerGluPheLeuLeuGlnLeuAspSerCysHisLeuAspLeuGlyProGluGlyGlyThrValGluLeuIle

1801 : CAGGGCGGGCGGCCAAGGGCAACTGTGTAGCCTGCTGTCCCCAAGCCCGAGGGTGACCCGCGCTTACGTTCTCTCTCCACTTCTAC
508 : GlnGlyArgAlaAlaLysGlyAsnCysValSerLeuLeuSerProSerProGluGlyAspProArgPheSerPheLeuLeuHisPheTyr

1891 : ACAGTACCATACCCAAACCGGCCACTCTCAGCTGCACGGTAGCCCTCGCTCCCAAGACCGGGTCTCAAGACCAGGAAGTCCATAGGACT
538 : ThrValProIleProLeuSerThrGlyThrLeuSerCysThrValAlaLeuArgProLysThrGlySerGlnAspGlnGluValHisArgThr

1981 : GTCTCATGCGCTTGAACATCATCAGCCCTGACTGCTGTGGTTGCACAAGAAGCCCTCTGCTGCCCGCGTGTGGGCATCACTT
568 : ValPheMetArgLeuAsnIleIleSerProAspLeuSerGlyCysThrSerLysGlyLeuValLeuProAlaValGlyIleThrPhe

2071 : GGTGCTTCTCTCATCGGGCCCTGCTCACTGCTGCACTTGGTACATCTACTCGCACAGCGGTTCCCCCAGCAAGCGGGAGCCGCTGCTG
598 : GlyAlaPheLeuIleGlyAlaLeuLeuThrAlaAlaLeuTrpTyrIleTyrSerHisThrArgSerProSerLysArgGluProValVal

2161 : CGGTGGCTGCCCGGCCCTCTCGGAGAGCAGCAGCCACACAGCATCGGAGCACCCAGAGCACCCCTGCTCCACCAGCAGCATG
628 : AlaValAlaAlaProAlaSerSerGluSerSerSerThrAsnHisSerIleGlySerThrGlnSerThrProCysSerThrSerSerMet

2251 : GCATGCCCGCGCCCCCGGCTCGCCCGAGCAGGAGACTGAGCAGCGCCAGCTGGGAGCACTGGTGTGAACCTCACCTGGGAGCCAG
658 : Alastop.....
2341 : TCTCTCACTCGACCAGAAATGGAGCGTGCTCTCGCCGCTTACCTCTCCCGCTCTCTCTCGAGAGCCTGCTGCCAGTGCAGCCACTGCGCT
2431 : TGGAAACACTTGGGCTCTCTCAACCCACAGAACTTCAACCACTGGGTGTGGAGTATGGCTCCCAAGGAGACAGACCTTGGCAGCC
2521 : TGTGTAAAAACCAAGTCTCTGATTTGAACCTGGATCCAGCACTGGTGAACCTGAGCTGGGAGGAGGGGAGCCAGCACTTGATCTATCTCTCGGCCCTC
2611 : AGGCCAACCCAGCCAGGACCCAGCAGCTCTCCGCTCGGACAGAGAGGGGGCCGACCCAGGACCACTGGATCTATCTCTCGGCCCTC
2701 : ACACCTGTAACCTGCTTAACTAACTGCGACGGACAGCAGAGCTAGCCAGGCTAGCCCTAGCCCTGGGAGCCAGGAGGTGGCAAGAACAGCTGGGG
2791 : TCTGGAGACTAGCTCTGCCACATATGAGCCCTCTCGGCTCGGGCAGCCAGCAGAGGGGGAGTAGCCCAAGCTGCTGTGCTCGGCCCT
2881 : GCGCCCTGTGTATTACACCAATAAATGACAGCACTGAACCACTGAAAAAAAAAAAA

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US93/10307

A. CLASSIFICATION OF SUBJECT MATTER IPC(5) : Please See Extra Sheet. US CL : 435/70.1, 240.2, 252.3, 320.1; 514/2; 530/395; 536/23.5 According to International Patent Classification (IPC) or to both national classification and IPC																				
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 435/70.1, 240.2, 252.3, 320.1; 514/2; 530/395; 536/23.5 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) APS, Medline, Biosis search terms: endoglin, TGF-B																				
C. DOCUMENTS CONSIDERED TO BE RELEVANT																				
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.																		
X — Y — A	J. Biol. Chem., Vol. 265, Number 15, issued 25 May 1990, Gougos et al., "Primary structure of endoglin, an RGD-containing glycoprotein of human endothelial cells", pages 8361-8364, see Fig. 4.	1-4 — 5-12 — 13-46																		
X — A	J. Biol. Chem., Vol. 267, Number 27, issued 25 September 1992, Cheifetz et al., "Endoglin is a component of the transforming growth factor-Beta receptor system in human endothelial cells", pages 19027-19030, see Fig. 4.	6, 10-12 — 13-46																		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.																				
<table border="0"> <tr> <td>* Special categories of cited documents:</td> <td>T</td> <td>later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</td> </tr> <tr> <td>*A* document defining the general state of the art which is not considered to be part of particular relevance</td> <td>X*</td> <td>document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td> </tr> <tr> <td>*E* earlier document published on or after the international filing date</td> <td>Y*</td> <td>document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</td> </tr> <tr> <td>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reasons (as specified)</td> <td>A*</td> <td>document member of the same patent family</td> </tr> <tr> <td>*O* document referring to an oral disclosure, use, exhibition or other means</td> <td></td> <td></td> </tr> <tr> <td>*P* document published prior to the international filing date but later than the priority date claimed</td> <td></td> <td></td> </tr> </table>			* Special categories of cited documents:	T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	*A* document defining the general state of the art which is not considered to be part of particular relevance	X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	*E* earlier document published on or after the international filing date	Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reasons (as specified)	A*	document member of the same patent family	*O* document referring to an oral disclosure, use, exhibition or other means			*P* document published prior to the international filing date but later than the priority date claimed		
* Special categories of cited documents:	T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention																		
A document defining the general state of the art which is not considered to be part of particular relevance	X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone																		
E earlier document published on or after the international filing date	Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art																		
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reasons (as specified)	A*	document member of the same patent family																		
O document referring to an oral disclosure, use, exhibition or other means																				
P document published prior to the international filing date but later than the priority date claimed																				
Date of the actual completion of the international search 12 January 1994		Date of mailing of the international search report 02 FEB 1994																		
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231		Authorized officer KENNETH R. HORLICK <i>K. R. Horlick</i>																		
Facsimile No. NOT APPLICABLE		Telephone No. (703) 305-6126																		

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US93/10307

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	J. Immunol., Vol. 141, Number 6, issued 15 September 1988, Gougos et al., "Identification of a human endothelial cell antigen with monoclonal antibody 44G4 produced against a pre-B leukemic cell line", pages 1925-1933, see Fig. 8.	11-12

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US93/10307

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (5):

C07H 21/02, 04; C07K 9/00; A61K 37/00; C12N 1/21, 5/10, 15/70; C12P 21/00

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☒ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☒ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☒ **GRAY SCALE DOCUMENTS**

☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.